

Social Anxiety
Panic disorder and agoraphobia
Anxiety Disorder
Obsessive Compulsive Spectrum Disorder
Social Anxiety
Specific Phobias

CBT FOR ANXIETY DISORDERS

A Practitioner Book

EDITED BY

GREGORIS SIMOS

AND STEFAN G. HOFMANN

 WILEY-BLACKWELL

CBT for Anxiety Disorders

CBT for Anxiety Disorders

A Practitioner Book

Edited by

Gregoris Simos

University of Macedonia, Thessaloniki, Greece

Stefan G. Hofmann

Boston University, Boston, MA, USA

 **WILEY-BLACKWELL**

A John Wiley & Sons, Ltd., Publication

This edition first published 2013
© 2013 John Wiley & Sons, Ltd.

Blackwell Publishing was acquired by John Wiley & Sons in February 2007. Blackwell's publishing program has been merged with Wiley's global Scientific, Technical, and Medical business to form Wiley-Blackwell.

Registered Office

John Wiley & Sons Ltd, The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK

Editorial Offices

350 Main Street, Malden, MA 02148-5020, USA

9600 Garsington Road, Oxford, OX4 2DQ, UK

The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK

For details of our global editorial offices, for customer services, and for information about how to apply for permission to reuse the copyright material in this book please see our website at www.wiley.com/wiley-blackwell.

The right of Gregoris Simos and Stefan G. Hofmann to be identified as the authors of this work/of the editorial material in this work has been asserted in accordance with the UK Copyright, Designs and Patents Act 1988.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, except as permitted by the UK Copyright, Designs and Patents Act 1988, without the prior permission of the publisher.

Wiley also publishes its books in a variety of electronic formats. Some content that appears in print may not be available in electronic books.

Designations used by companies to distinguish their products are often claimed as trademarks. All brand names and product names used in this book are trade names, service marks, trademarks or registered trademarks of their respective owners. The publisher is not associated with any product or vendor mentioned in this book. This publication is designed to provide accurate and authoritative information in regard to the subject matter covered. It is sold on the understanding that the publisher is not engaged in rendering professional services. If professional advice or other expert assistance is required, the services of a competent professional should be sought.

Library of Congress Cataloging-in-Publication Data

CBT for anxiety disorders : a practitioner book / edited by Gregoris Simos and Stefan G. Hofmann.
pages cm

Includes bibliographical references.

ISBN 978-0-470-97552-7 (cloth) – ISBN 978-0-470-97553-4 (pbk.) 1. Anxiety disorders – Treatment. 2. Cognitive therapy. I. Simos, Gregoris, 1953– editor of compilation. II. Hofmann, Stefan G., editor of compilation. III. Title: Cognitive behavior therapy for anxiety disorders.

RC531.C42 2013

616.85'22–dc23

2012047737

A catalogue record for this book is available from the British Library.

Cover design by Cyan Design

Typeset in 10/12.5pt Galliard by Aptara Inc., New Delhi, India

Contents

List of Contributors	vii
List of Figures, Tables, and Boxes	ix
Introduction	1
<i>Gregoris Simos and Stefan G. Hofmann</i>	
1 Panic Disorder and Agoraphobia	3
<i>Michelle G. Craske and Gregoris Simos</i>	
2 Obsessive–Compulsive Spectrum Disorders: Diagnosis, Theory, and Treatment	25
<i>David A. Clark and Gregoris Simos</i>	
3 Generalized Anxiety Disorder: Targeting Intolerance of Uncertainty	57
<i>Melisa Robichaud</i>	
4 Social Anxiety Disorder: Treatment Targets and Strategies	87
<i>Stefan G. Hofmann, Jacqueline Bullis, and Cassidy Gutner</i>	
5 Specific Phobias	107
<i>Lars-Göran Öst and Lena Reuterskiöld</i>	
6 Health Anxiety	135
<i>Michel A. Thibodeau, Gordon J.G. Asmundson, and Steven Taylor</i>	
7 Trauma-Focused Cognitive Behavior Therapy for Posttraumatic Stress Disorder and Acute Stress Disorder	161
<i>Anke Ehlers</i>	
8 Culturally Appropriate CBT for the Anxiety Disorders	191
<i>Devon E. Hinton and Martin La Roche</i>	
9 Newer Generations of CBT for Anxiety Disorders	225
<i>Michael P. Twohig, Michelle R. Woidneck, and Jesse M. Crosby</i>	
Index	251

List of Contributors

Gordon J.G. Asmundson, Department of Psychology, University of Regina, Regina, Canada

Jacqueline Bullis, Department of Psychology, Boston University, Boston, MA, USA

David A. Clark, Department of Psychology, University of New Brunswick, Fredericton, Canada

Michelle G. Craske, Department of Psychology, UCLA, Los Angeles, CA, USA

Jesse M. Crosby, McLean Hospital/Harvard Medical School, Belmont, MA, USA

Anke Ehlers, Department of Experimental Psychology, University of Oxford, Oxford, UK

Cassidy Gutner, Department of Psychology, Boston University, Boston, MA, USA

Devon E. Hinton, Department of Psychiatry, Massachusetts General Hospital, Boston, MA, USA; Arbour Counseling Services, Lowell, MA, USA; Children's Hospital/Martha Eliot Health Center and Harvard Medical School, Jamaica Plain, MA, USA

Stefan G. Hofmann, Department of Psychology, Boston University, Boston, MA, USA

Lars-Göran Öst, Department of Psychology, Stockholm University, Stockholm, Sweden

Lena Reuterskiöld, Department of Psychology, Stockholm University, Stockholm, Sweden

Melisa Robichaud, Vancouver CBT Centre, Vancouver, Canada; University of British Columbia, Department of Psychiatry, Vancouver, Canada

Martin La Roche, Children's Hospital/Martha Eliot Health Center and Harvard Medical School, Jamaica Plain, MA, USA

Gregoris Simos, Department of Educational and Social Policy, University of Macedonia, Thessaloniki, Greece

Steven Taylor, Department of Psychiatry, University of British Columbia, Vancouver, Canada

Michel A. Thibodeau, Department of Psychology, University of Regina, Regina, Canada

Michael P. Twohig, Department of Psychology, Utah State University, Logan, UT, USA

Michelle R. Woidneck, Department of Psychology, Utah State University, Logan, UT, USA

List of Figures, Tables, and Boxes

Figure 3.1	Illustrative GAD model for cognitive behavioral treatment targeting IU.	64
Table 3.1	Session-by-session treatment plan.	63
Table 3.2	Sample worry monitoring form.	65
Table 3.3	Examples of IU safety-seeking behaviors.	67
Table 3.4	Sample behavioral experiment record form.	69
Table 3.5	Positive beliefs about worry: examples and challenges.	72
Figure 4.1	Positive feedback loop that maintains SAD. (Reprinted with permission from Hofmann, 2007.)	90
Figure 5.1	Daily self-ratings of anxiety during baseline and posttreatment periods (↓ = the day of each treatment session).	128
Table 5.1	Proposed DSM-V criteria for specific phobia.	108
Figure 6.1	Theoretical illustration of Robert's severe health anxiety and the environmental (solid rectangles), behavioral (solid circles), physiological (lined circles), and cognitive features (lined rectangles) that maintain his anxiety.	145
Figure 7.1	Treatment goals in Cognitive Therapy for PTSD. Pointed arrows stand for "leads to." Round arrows stand for "prevents a change in." Dashed arrows stand for "influences."	166
Box 7.1	Example of an updated trauma narrative (extracts, updates in CAPITALS).	171
Table 7.1	Stimulus discrimination: examples of sensory triggers of reexperiencing symptoms.	172

Table 7.2	Steps in behavioral experiments: example from the treatment of an assault survivor.	174
Figure 8.1	A culturally sensitive model of how anxiety disorders occur in a certain cultural context: psychopathological mechanisms.	192
Figure 8.2	A culturally sensitive model of how anxiety disorders occur in a certain cultural context: balance of psychopathological mechanisms and local response.	193
Figure 8.3	A culturally sensitive model of how CBT can reduce the severity of anxiety disorders.	198
Figure 8.4	The nodal network model of negative affect: focus on inflexibility aspects. This is a nodal network model of negative affect, showing how multiple nodes interact to create a negative affective state. At one point in time, one or another node may be the object of attention or may be more active in determining the current mode of processing. As one node becomes active, all the other nodes tend to be so as well; if one node shifts, all the others tend to as well.	217
Figure 8.5	The nodal network model of positive affect: focus on flexibility aspects. This is a nodal network model of positive affect, showing how multiple nodes interact to create a positive affective state. At one point in time, one or another node may be the object of attention or may be more active in determining the current mode of processing. As one node becomes active, all the other nodes tend to be so as well; if one node shifts, all the others tend to as well.	218
Table 8.1	Anxiety-related distress syndromes in Asia: associated catastrophic cognitions, DSM-IV diagnosis, and panic attack subtype.	194
Table 8.2	Sessions in CA-CBT and key components of the sessions.	199
Table 8.3	Ways to make CBT-type treatment of anxiety disorders for other cultural groups to be more efficacious and culturally appropriate.	200
Table 8.4	Examples of proverbs and culturally salient expressions that can be used therapeutically.	206
Figure 9.1	Illustration of the “umbrella of CBT.”	232

Introduction

Gregoris Simos¹ and Stefan G. Hofmann²

¹*University of Macedonia, Thessaloniki, Greece*

²*Boston University, Boston, MA, USA*

Anxiety disorders are some of the most prevalent disorders, affecting three out of ten people in their lifetime (Kessler *et al.*, 2005). These disorders start early in life, negatively affect significant aspects of functioning, tend to be chronic and unremitting, and cause considerable psychological suffering and life impairments (Martin, 2003). In addition, they are highly comorbid with three out of four anxiety disorder patients experience at least one other mental disorder in their lifetime (Michael, Zetsche, and Margraf, 2007).

The efficacy of cognitive behavior therapy (CBT) for anxiety disorders in adults has been supported by multiple meta-analyses (Hofmann *et al.*, 2012). For example, CBT is superior to no treatment and control treatments (Norton and Price, 2007; Olatunji, Cisler, and Deacon, 2010), including placebo control (Hofmann and Smits, 2008). The latter meta-analysis shows that CBT is associated with medium to large effect sizes over placebo, suggesting that, although effective, there is still considerable room for further improvement (Hofmann and Smits, 2008).

Research on the cognitive model of anxiety and the development of disorder-specific cognitive treatment protocols for anxiety disorders is a continuously evolving process (e.g., Clark and Beck, 2010). For this reason, we invited some of the foremost experts on CBT for anxiety disorders to provide an update of the contemporary state of the art of treating anxiety disorders. All chapters include the treatment rationale, concrete clinical case examples, therapist–patient dialogues and, where appropriate, subsections of techniques for dealing with treatment complications, comorbid disorders, and managing concurrent pharmacotherapy, and ethnicity.

The list of contributors and topics include “Panic Disorder and Agoraphobia” (Chapter 1) by Michelle G. Craske and Gregoris Simos; “Obsessive–Compulsive Spectrum Disorders: Diagnosis, Theory, and Treatment” (Chapter 2) by David A. Clark and Gregoris Simos; “Generalized Anxiety Disorder: Targeting Intolerance of Uncertainty” (Chapter 3) by Melisa Robichaud; “Social Anxiety Disorder: Treatment Targets and Strategies” (Chapter 4) by Stefan G. Hofmann, Jacqueline Bullis, and Cassidy

Gutner; “Specific Phobias” (Chapter 5) by Lars-Göran Öst and Lena Reuterskiöld; “Health Anxiety” (Chapter 6) by Michel A. Thibodeau, Gordon J.G. Asmundson, and Steven Taylor; “Trauma-Focused Cognitive Behavior Therapy for Posttraumatic Stress Disorder and Acute Stress Disorder” (Chapter 7) by Anke Ehlers; “Culturally Appropriate CBT for the Anxiety Disorders” (Chapter 8) by Devon E. Hinton and Martin La Roche; and “Newer Generations of CBT for Anxiety Disorders” (Chapter 9) by Michael P. Twohig, Michelle R. Woidneck, and Jesse M. Crosby.

Our hope is that this text will provide the reader with up-to-date knowledge about the current state-of-the-art CBT approaches for anxiety disorders. We believe that it will be of interest to anyone who wants to help patients with anxiety disorders – practitioners in training, senior clinicians, researchers, residents, graduate psychology, and medical students.

References

- Clark, D.A. and Beck, A.T. (2010) *Cognitive Therapy of Anxiety Disorders: Science and Practice*, Guilford Press, New York.
- Hofmann, S.G. and Smits, J.A. (2008) Cognitive-behavioral therapy for adult anxiety disorders: a meta-analysis of randomized placebo-controlled trials. *Journal of Clinical Psychiatry*, 69, 621–632.
- Hofmann, S.G., Asnaani, A., Vonk, I.J.J. *et al.* (2012) The efficacy of cognitive behavioral therapy: A review of meta-analyses. *Cognitive Therapy and Research*, 36, 427–440.
- Kessler, R.C., Berglund, P., Demler, O. *et al.* (2005) Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Archives of General Psychiatry*, 62, 593–602.
- Martin, P. (2003) The epidemiology of anxiety disorders: a review. *Dialogues in Clinical Neuroscience*, 5, 281–298.
- Michael, T., Zetsche, U., and Margraf, J. (2007) Epidemiology of anxiety disorders. *Psychiatry*, 6, 136–142.
- Norton, P.J. and Price, E.C. (2007) A meta-analytic review of adult cognitive-behavioral treatment outcome across the anxiety disorders. *Journal of Nervous and Mental Diseases*, 195, 521–531.
- Olatunji, B.O., Cisler, J.M., and Deacon, B.J. (2010) Efficacy of cognitive behavioral therapy for anxiety disorders: a review of meta-analytic findings. *Psychiatric Clinics of North America*, 33, 557–577.

Panic Disorder and Agoraphobia

Michelle G. Craske¹ and Gregoris Simos²

¹*UCLA, Los Angeles, CA, USA*

²*University of Macedonia, Thessaloniki, Greece*

Overview

The current diagnostic criteria for panic disorder, according to the *Diagnostic and Statistical Manual of Mental Disorders*, 4th edition (DSM-IV, American Psychiatric Association, 1994), are comprised of recurrent unexpected panic attacks, and anxiety about future panic attacks or their consequences, or a significant behavioral change because of the panic attacks. The additional anxiety about panic, combined with catastrophic cognitions about panic sensations, contributes to the differentiation between the person with panic disorder and the person with occasional panic attacks (e.g., Telch, Lucas, and Nelson, 1989). Agoraphobia refers to avoidance or endurance with dread of situations from which escape might be difficult or help unavailable in the event of a panic attack, or in the event of developing symptoms that could be incapacitating and embarrassing, such as loss of bowel control or vomiting. Typical agoraphobic situations include shopping malls, waiting in line, movie theaters, traveling by car or bus, crowded restaurants, and being alone.

In the general population, the 12-month prevalence estimate for panic disorder across the United States and several European countries is about 2% in adults and adolescents (Goodwin, Fergusson, and Horwood, 2005; Kessler *et al.*, 2005b). Lower estimates have been reported for some Asian, African, and Latin American countries, ranging from 0.1 to 0.8% (Lewis-Fernandez *et al.*, 2010). Across all studies, females are more frequently affected than males at a rate of approximately 2 : 1 (Kessler *et al.*, 2005b). Although panic attacks occur in children, the overall prevalence of panic disorder is low prior to 14 years of age (<0.4%) (Craske *et al.*, 2010). The rates of panic disorder show a gradual increase during adolescence, particularly in girls, and possibly following the onset of puberty (Craske *et al.*, 2010). The modal age of onset is late teenage years and early adulthood (Kessler *et al.*, 2005a), although treatment is usually sought at a much later age, around 34 years (e.g., Noyes *et al.*, 1986). The

prevalence rates decline in older individuals, possibly reflecting diminishing severity to subclinical levels (Wolitzky-Taylor *et al.*, 2010). In general, differences in prevalence across gender, culture, and age groups may be due to a variety of factors, including the expression of the disorder, underlying physiology or biology, varying degrees of concern about the dangerousness of symptoms of autonomic arousal and mental symptoms of anxiety, and sensitivity of instrumentation for diagnosing panic disorder.

Panic disorder and agoraphobia tend to be chronic conditions, with severe financial and interpersonal costs; that is, only a minority (30%) of untreated individuals remit without subsequent relapse, although a similar number experience notable improvement, albeit with a waxing and waning course (35%) (Katschnig and Amering, 1998; Roy-Byrne and Cowley, 1995). Also, panic disorder is associated with high levels of social, occupational, and physical disability, considerable economic costs, and the highest number of medical visits among the anxiety disorders, although the effects are strongest with the presence of agoraphobia (Wittchen *et al.*, 2010).

Rarely does the diagnosis of panic disorder, with or without agoraphobia, occur in isolation. Commonly co-occurring Axis I conditions include specific phobias, social phobia, dysthymia, generalized anxiety disorder, major depressive disorder, and substance abuse (e.g., Brown *et al.*, 2001; Kessler *et al.*, 2005b). Also, from 25 to 60% of persons with panic disorder also meet criteria for a personality disorder, mostly avoidant and dependent personality disorders (e.g., Chambless and Renneberg, 1988). However, the nature of the relationship between panic disorder/agoraphobia and personality disorders remains unclear, especially as some “personality disorders” remit after successful treatment of panic disorder/agoraphobia (e.g., Latas *et al.*, 2000; Marchesi *et al.*, 2005; Ozkan and Altindag, 2005).

Cognitive Behavioral Model

Several independent lines of research (Barlow, 1988; Clark, 1986; Ehlers and Margraf, 1989) converged in the 1980s on the same basic conceptualization of panic disorder as an acquired fear of bodily sensations, particularly sensations associated with autonomic arousal. Psychological (i.e., temperament, such as negative affectivity) and biological (i.e., genetic) predispositions are believed to enhance the vulnerability to acquire such fear. Fear conditioning, avoidant responding, and information processing biases are believed to perpetuate such fear. It is the perpetuating factors that are targeted in the cognitive behavioral treatment approach.

The temperament most associated with anxiety disorders, including panic disorder, is neuroticism (Eysenck, 1967; Gray, 1982) or proneness to experience negative emotions in response to stressors. A closely linked construct is “negative affect,” or the tendency to experience a variety of negative emotions across a variety of situations, even in the absence of objective stressors (Watson and Clark, 1984). Neuroticism predicts the onset of panic attacks in adolescents (Hayward *et al.*, 2000; Schmidt, Lerew and Jackson, 1997, 1999), and “emotional reactivity” at age 3 was a significant variable in the classification of panic disorder in 18- to 21-year-old males (Craske *et al.*, 2001). Numerous multivariate genetic analyses of human twin samples consistently

attribute approximately 30–50% of variance in neuroticism to additive genetic factors (Eley, 2001; Lake *et al.*, 2000). In addition, anxiety and depression appear to be variable expressions of the heritable tendency toward neuroticism (Kendler *et al.*, 1987). Symptoms of panic (i.e., breathlessness and heart pounding) may be additionally explained by a unique source of genetic variance that is differentiated from symptoms of depression and anxiety (Kendler *et al.*, 1987) and neuroticism (Martin *et al.*, 1988).

Another temperament is anxiety sensitivity, which refers to the trait of believing that anxiety and associated symptoms may cause deleterious physical, social, and psychological consequences that extend beyond any immediate physical discomfort during an episode of anxiety or panic (Reiss, 1980). Anxiety sensitivity is elevated across most anxiety disorders, but it is particularly elevated in panic disorder, especially the physical concerns subscale (Zinbarg, Barlow, and Brown, 1997). Anxiety sensitivity is believed to comprise a specific psychological vulnerability for panic disorder because it primes fear reactivity to bodily sensations. In support, several longitudinal studies indicate that high scores on the anxiety sensitivity index predict the onset of panic attacks over 1- to 4-year intervals in adolescents (Hayward *et al.*, 2000), college students (Maller and Reiss, 1992), and community samples with specific phobias or no anxiety disorders (Ehlers, 1995). In addition, anxiety sensitivity index scores predicted spontaneous panic attacks, and worry about panic (and anxiety more generally), during an acute military stressor (i.e., 5 weeks of basic training), even after controlling for history of panic attacks and trait anxiety (Schmidt, Lerew and Jackson, 1999). Finally, panic attacks themselves elevate anxiety sensitivity over a 5-week period in adults (Schmidt Lerew, and Jackson, 1999), and over a 1-year period in adolescents, albeit to a lesser extent (Weems *et al.*, 2002).

However, Bouton, Mineka, and Barlow (2001) noted that the relationship between anxiety sensitivity and panic attacks in these studies was relatively small, not exclusive to panic, and was weaker than the relationship between panic and neuroticism. Furthermore, these studies have evaluated panic attacks and worry about panic but have not evaluated prediction of diagnosed panic disorder. Thus, the causal significance of anxiety sensitivity for panic disorder remains to be fully understood.

Acute “fear of fear” (or more accurately, anxiety focused on somatic sensations) that develops after initial panic attacks is attributed to two factors: catastrophic misappraisals of bodily sensations (i.e., misinterpretation of sensations as signs of imminent death, loss of control, and so on) (Clark, 1986); and interoceptive conditioning, or conditioned fear of internal cues, such as elevated heart rate, because of their association with intense fear, pain, or distress (Razran, 1961). Specifically, interoceptive conditioning refers to low-level bodily sensations of arousal coming to serve as conditional stimuli that trigger increased autonomic arousal and fear through Pavlovian conditioning (Bouton, Mineka, and Barlow, 2001). Thus, small changes in physiological functioning lead to conditioned fear or panic as a result of prior pairings of these initial somatic sensations with full-blown panic attacks. An extensive experimental literature attests to the robustness of interoceptive conditioning (e.g., Dworkin and Dworkin, 1999), particularly with regard to early interoceptive drug onset cues becoming conditioned stimuli for larger drug effects (e.g., Sokolowska, Siegel, and Kim, 2002). In addition, interoceptive conditioned responses are not dependent on

conscious awareness of triggering cues and thus have been observed under anesthesia (e.g., Block *et al.*, 1987). As such, interoceptive conditioning accounts for what appear to be “out of the blue” panic attacks.

Evidence for extreme fear and anxiety of somatic sensations is robust across a variety of paradigms. Persons with panic disorder endorse strong beliefs that bodily sensations associated with panic attacks cause physical or mental harm (e.g., McNally and Lorenz, 1987). They are more likely to interpret bodily sensations in a catastrophic fashion (Clark *et al.*, 1988), and to allocate more attentional resources to words that represent physical threat such as “disease” and “fatality” (e.g., Hope *et al.*, 1990), catastrophe words, such as “death” and “insane” (e.g., Maidenberg *et al.*, 1996), and heartbeat stimuli (Kroeze and van den Hout, 2000). Also, individuals with panic disorder show enhanced brain potentials to panic-related words (Pauli *et al.*, 2005). In addition, they are more likely to become anxious in procedures that elicit bodily sensations similar to the ones experienced during panic attacks, including benign cardiovascular, respiratory, and audiovestibular exercises (Antony *et al.*, 2006), as well as more invasive procedures such as carbon dioxide inhalations, compared to clients with other anxiety disorders (e.g., Perna *et al.*, 1995; Rapee *et al.*, 1992) or healthy controls (e.g., Gorman *et al.*, 1994). The findings are not fully consistent, however, as clients with panic disorder did not differ from clients with social phobia in response to an epinephrine challenge (Veltman *et al.*, 1996). Nonetheless, individuals with panic disorder also fear signals that ostensibly reflect heightened arousal and false physiological feedback (Craske *et al.*, 2002; Ehlers *et al.*, 1988).

Such anxiety about bodily sensations plays a central role in the perpetuation of panic disorder. First, once the bodily sensations are noticed, they elicit fear in an individual with panic disorder. This fear serves to intensify the sensations, causing an increase in fear, which further enhances the bodily sensations in a self-perpetuating cycle of fear and bodily sensations that typically results in a panic attack. Second, because bodily sensations that trigger panic attacks are not always immediately obvious, they may generate the perception of unexpected or “out of the blue” panic attacks that generates even further distress (Craske, Glover, and DeCola, 1995). Third, the perceived uncontrollability, or inability to escape from, or terminate bodily sensations again is likely to generate heightened anxiety (e.g., Maier, Laudenslager, and Ryan, 1985). Unpredictability and uncontrollability, then, are seen as enhancing general levels of anxiety about “when is it going to happen again” and “what do I do when it happens,” thereby contributing to high levels of chronic anxious apprehension. In turn, anxious apprehension increases the likelihood of panic, by directly increasing the availability of sensations that have become conditioned cues for panic and/or by increasing attentional vigilance for these bodily cues. Thus, a maintaining cycle of panic and anxious apprehension develops.

Individuals with panic disorder often engage in safety behaviors that they believe enable them to escape or avoid the feared outcome. For example, if individuals believe that they will pass out during a panic attack, they might sit down or hold on to an object for support. Engaging in safety behaviors prevents disconfirmation of cognitive misappraisals, thus contributing to the maintenance of panic disorder (Salkovskis, Clark, and Gelder, 1996). Individuals may also engage in safety behaviors designed to prevent panic, or its feared consequences, such as carrying around anxiolytic medication

or traveling with a companion who makes them feel safe. Another panic-maintaining behavioral response is overt avoidance of particular places or situations where panic attacks are anticipated to occur. Avoidance prevents disconfirmation of catastrophic misappraisals, and reinforces the fear that those particular situations are dangerous, increasing the likelihood of panicking in those situations in the future.

Components of Cognitive Behavioral Therapy

As outlined in more detail elsewhere (Simos, 2002), the treatment begins with education about the nature of panic, the causes of panic and anxiety, and the way in which panic and anxiety are perpetuated by feedback loops among physical, cognitive, and behavioral response systems. In addition, specific descriptions of the psychophysiology of the fight-flight response are provided, as well as an explanation of the adaptive value of the various physiological changes that occur during panic and anxiety. The purpose of this education is to correct common myths and misconceptions about panic symptoms (i.e., beliefs about going crazy, dying, or losing control).

Self-monitoring is introduced in the first treatment session and is continued throughout the entire treatment. Self-monitoring functions in two ways: to provide ongoing assessment of change in panic, anxiety, and avoidance; and as a therapeutic tool to encourage objective self-awareness and increase accuracy in self-observation. Clients are asked to keep at least two types of self-monitoring records. The first is a panic attack record, to be completed as soon as possible after each panic attack; this record provides a description of cues, maximal distress, symptoms, thoughts, and behaviors. The second is a daily mood record, completed at the end of each day, to keep record of overall or average levels of anxiety, depression, and so on. Additionally, clients may keep a daily record of activities or situations avoided.

Breathing retraining is a commonly used somatic coping skill, given evidence for respiratory abnormalities in panic disorder possibly due to hypersensitive medullary carbon dioxide (CO_2) detectors, resulting in hypocapnia (i.e., lower than normal levels of $p\text{CO}_2$) (e.g., Caldirola *et al.*, 2004). Traditional breathing retraining involves slow, abdominal breathing exercises. However, its value has been questioned in terms of the degree to which it actually corrects hypocapnic breathing or rather serves as a distraction (Garssen, de Ruiter, and van Dyck, 1992). In contrast to traditional breathing retraining, capnometry-assisted respiratory training (CART) (Meuret *et al.*, 2008) uses immediate feedback of end-tidal $p\text{CO}_2$ to teach clients how to raise their subnormal levels of $p\text{CO}_2$ (hyperventilation) and thereby gain control over dysfunctional respiratory patterns and associated panic symptoms (e.g., shortness of breath and dizziness). CART has been shown to improve panic symptoms, in part through reducing hypocapnic breathing (Meuret *et al.*, 2010). Another somatic coping skill is progressive muscle relaxation, in which clients are trained over a number of weeks in 16-muscle groups, 8-muscle groups, 4-muscle groups, and finally cue-control relaxation, at which point relaxation is used as a coping skill for practicing exposure to items from a hierarchy of anxiety-provoking tasks.

In the cognitive restructuring component of cognitive behavioral therapy (CBT), detailed self-monitoring of emotions and associated cognitions is used to identify

specific beliefs, appraisals, and assumptions. Relevant cognitions are categorized into types of errors, such as overestimations of risk of negative events, or catastrophizing the meaning of events. In labeling the type of cognitive distortion, the client is encouraged to use an empirical approach to examine the validity of thoughts by considering all of the available evidence. Therapists use Socratic questioning to help clients make guided discoveries and question their anxious thoughts. Next, alternative hypotheses are generated that are more evidence-based. In addition to surface-level appraisals (such as “my heart is racing dangerously too fast”), core-level beliefs or schemata (such as “I am too weak to withstand distress”) are questioned in the same way.

In vivo exposure refers to repeated and systematic real-life exposure to agoraphobic situations. Most often, *in vivo* exposure is conducted in a graduated manner, proceeding from the least to the most anxiety-provoking situations on an avoidance hierarchy, although there is some evidence to suggest that intensive or ungraduated exposure may be effective (e.g., Feigenbaum, 1988). Critical to *in vivo* exposure is the removal of safety signals and safety behaviors, such as other people, empty or full medication bottles, seeking reassurance, or checking for exits. Reliance on safety signals and safety behaviors attenuates distress in the short term but is believed to maintain excessive anxiety in the long term. They are replaced by effective use of cognitive restructuring and somatic coping skills, with care to ensure that the coping skills themselves do not become alternative safety behaviors. *In vivo* exposure can be conducted with the therapist’s guidance, followed by self-directed exposures between sessions (to enhance generalization of learning and to limit the safety signal value of the therapist). Recent data support the value of therapist-directed exposure (Gloster *et al.*, 2011).

In interoceptive exposure, the goal is to deliberately induce feared physical sensations a sufficient number of times and for long enough each time so that misappraisals about the sensations are disconfirmed and conditioned anxiety responding extinguishes. A standard list of exercises, such as hyperventilating and spinning, are used to establish a hierarchy of interoceptive exposures. Clients are encouraged to endure the sensations beyond the point at which they are first noticed because early termination interferes with new learning. Interoceptive exposure is usually first conducted in-session with the therapist’s guidance, followed by self-directed practice between sessions. Interoceptive exposure extends to naturalistic activities that inherently induce somatic sensations (e.g., caffeine consumption and exercise programs). Eventually, *in vivo* exposure is combined with interoceptive exposure, by deliberately inducing feared sensations in feared situations.

A final step of CBT is relapse prevention, in which clients are informed that recurrences of panic, anxiety or avoidance behavior are likely to occur in the future. They are encouraged to view such recurrences as lapses rather than failure, and to reapply their coping skills and reinstitute their practice of interoceptive and *in vivo* exposure.

Science of Exposure Therapy

Exposure therapy has developed over time, originating with graduated imaginal exposure combined with counterconditioning through relaxation (i.e., systematic desensitization) developed by Wolpe (1959). Emotional processing theory emphasized

habituation of fear responding within an exposure trial as a necessary precursor to habituation across treatment sessions, with the aim of long-term corrective learning (Foa and Kozak, 1986; Foa and McNally, 1996). Most recently, we have emphasized optimizing inhibitory learning and its retrieval in ways that are not necessarily dependent on reductions in fear throughout trials of exposure (Craske *et al.*, 2008); we discuss this approach below.

Emotional processing theory emphasizes mechanisms of habituation as precursors to cognitive correction. Specifically, emotional processing theory purports that the effects of exposure therapy derive from activation of a “fear structure” and integration of information that is incompatible with it, resulting in the development of a nonfear structure that replaces or competes with the original one. Incompatible information derives first from within-session habituation, or reduction in fear responding with prolonged exposure to the fear stimulus. Within-session habituation is seen as a prerequisite for the second piece of incompatible information, which derives from between-session habituation over repeated occasions of exposure. Between-session habituation is purported to form the basis for long-term learning and to be mediated by changes in “meaning,” or lowered probability of harm (i.e., risk) and lessened negativity (i.e., valence) of the stimulus. Emotional processing theory guides clinicians to focus on the initial elevation of fear followed by within- and between-session reductions in fear as signs of treatment success. Although enticing in its face validity, support for the theory has been inconsistent at best (Craske *et al.*, 2008, 2012). Rather, the evidence suggests that the amount by which fear habituates from the beginning to the end of an exposure practice is not a good predictor of overall outcomes, and that evidence for between-session habituation is mixed (Craske *et al.*, 2008, 2012).

A return to the science of fear learning and extinction may help to explain the effects of exposure therapy and thereby optimize its implementation. It is now thought that inhibitory learning is central to extinction (Bouton, 1993). Inhibitory pathways are also recognized in the neurobiology of fear extinction (see Sotres-Bayon, Cain, and LeDoux, 2006). Within a Pavlovian conditioning approach, inhibitory learning means that the original association between the conditional stimulus (i.e., the neutral stimulus that is paired with an innately aversive stimulus) and the unconditional stimulus (the innately aversive stimulus) learned during fear conditioning is not erased during extinction, but rather is left intact as a new, secondary learning (i.e., the conditional stimulus no longer predicts the unconditional stimulus) develops (Bouton, 1993). The degree to which inhibitory associations shape fear responding at retest (the index of strength and stability of new “learning”) is independent of fear levels expressed throughout extinction and instead is dependent on factors such as context and time.

Based on the inhibitory retrieval model of extinction, outcomes may be enhanced by strategies that do not rely on fear reduction within a trial of exposure (Craske *et al.*, 2008, 2012). Indeed, fear reduction may become a safety behavior for persons with panic disorder (since fear reduction eradicates the very thing that is feared), such that a more appropriate goal may be to maintain high levels of fear and anxiety in order to disconfirm the expectancy of negative consequences. One translational possibility is “deepened extinction” (Rescorla, 2006), where multiple fear conditional stimuli are first extinguished separately before being combined during extinction, and in animal studies, decreases spontaneous recovery and reinstatement of fear. Indeed, this is what

is essentially done when interoceptive exposure is conducted in feared agoraphobic situations (Barlow and Craske, 1994) and recent experimental data support the beneficial effects of deepened extinction in human conditioning studies (N.C. Culver, B. Vervliet, and M.G. Craske, manuscript in preparation).

In addition, the effects of exposure therapy may be enhanced by the prevention or removal of “safety signals” or “safety behaviors.” Common safety signals and behaviors for clients with panic disorder are the presence of another person, therapists, medications, or food or drink. In the experimental literature, safety signals alleviate distress in the short term, but when they are no longer present, the fear returns (Lovibond, Davis, and O’Flaherty, 2000), an effect that may derive in part from interference with the development of inhibitory associations. In phobic samples, the availability and use of safety signals and behaviors has been shown to be detrimental to exposure therapy (Sloan and Telch, 2002), whereas instructions to refrain from using safety behaviors improved outcomes (Salkovskis, 1991), although recent data have presented contradictory findings (Rachman *et al.*, 2011).

Further options include stimulus variability throughout exposure since variability has been shown to enhance the storage capacity of newly learned information. Two studies with clinical analogs have demonstrated positive benefits in terms of spontaneous recovery (Lang and Craske, 2000; Rowe and Craske, 1998), while a third showed trends only (Kircanski *et al.*, 2012). In the treatment for panic disorder/agoraphobia, this implies conducting exposure for varying durations, at varying levels of intensity, rather than continuing exposure in one situation until fear declines before moving to the next situation. Notably, such variability typically elicits higher levels of anxiety during exposure, but without detrimental effects and sometimes with beneficial effects in the long term.

Based on evidence for fear extinction to be weakened by antagonists of the glutamate receptors in the amygdala, Walker *et al.* (2002) tested and demonstrated that drug agonists of the same receptors, and in particular, D-cycloserine, enhanced extinction in animal studies. In a meta-analysis of the efficacy of D-cycloserine for anxiety disorders, Norberg, Krystal, and Tolin (2008) reported effect sizes of $d = 0.60$ at posttreatment and 0.47 at follow-up in clinical anxiety samples. D-Cycloserine in combination with interoceptive exposure for panic clients has resulted in a greater reduction in symptom severity, and a greater likelihood of achieving a change in clinical status at posttreatment and 1-month follow-up compared to exposure plus placebo (Otto *et al.*, 2010). Notably, D-cycloserine has been shown to have positive effects without influencing the level of fear during exposure per se.

A number of options for enhancing retrieval of the extinction memory have been tested. One option is to include retrieval cues during extinction training to be used in other contexts once extinction is over. This has been shown to be effective in animal studies and human conditioning studies (see Craske *et al.*, 2012, for a review). In clinical analog samples, the effects of a retrieval cue upon context renewal were very weak in one study (Culver, Stoyanova, and Craske, 2011), although instructions to mentally reinstate what was learned during exposure had more robust effects in reducing context renewal in another study (Mystkowski *et al.*, 2006). In the treatment of panic disorder, this approach simply suggests that clients carry cues (e.g., wrist band) with them to remind them of what they learned during exposure therapy (as long as

the cues do not become safety signals), or are prompted to remind themselves of what they learned in exposure therapy each time they experience previously feared sensations or situations.

Another option is to provide multiple contexts in which extinction takes place. This approach has been shown to offset context renewal in rodent samples, and in a clinical analog study of exposure therapy (Vansteenwegen *et al.*, 2007), although the results are not always consistent (Neumann, Lipp, and Cory, 2007). In the treatment of panic disorder and agoraphobia, this would mean asking clients to conduct their interoceptive and *in vivo* exposures in multiple different contexts, such as when alone, in unfamiliar places, or at varying times of the day or varying days of the week.

A recent (re-)discovery is that retrieving already stored memories induces a process of reconsolidation (Nader, Schafe, and Le Doux, 2000), since the memory is written into long-term memory again, requiring *de novo* neurochemical processes. Thus, it may be possible to change memories during the reconsolidation time frame upon retrieval. Propranolol, a β -blocker, has been shown to block the reconsolidation of memories, and Debiec and Ledoux (2004) found that infusions of propranolol blocked the reconsolidation of a previously formed CS-US memory, and led to erasure of the fear response and resistance to reinstatement effects. This suggests that propranolol upon retrieval may be a useful clinical tool, and indeed, two fear conditioning studies in healthy humans (Kindt, Soeter, and Vervliet, 2009; Soeter and Kindt, 2010) have replicated the effects. However, the effects have not been tested in the context of exposure therapy for panic disorder.

Differential Diagnosis

Panic disorder is not diagnosed if the panic attacks are judged to be a direct physiological consequence of a general medical condition, in which case an anxiety disorder due to a general medical condition is diagnosed. Examples of general medical conditions that can cause panic attacks include hyperthyroidism, hyperparathyroidism, pheochromocytoma, vestibular dysfunctions, seizure disorders, and cardiopulmonary conditions (e.g., arrhythmias, supraventricular tachycardia, asthma, and chronic obstructive pulmonary disease) (Fava *et al.*, 2010). Appropriate laboratory tests (e.g., serum calcium levels for hyperparathyroidism) or physical examinations (e.g., for cardiac conditions) may be helpful in determining the etiological role of a general medical condition.

Panic disorder is not diagnosed if the panic attacks are judged to be a direct physiological consequence of a substance (i.e., a drug of abuse and a medication), in which case a substance-induced anxiety disorder is diagnosed. Intoxication with central nervous system stimulants (e.g., cocaine, amphetamines, and caffeine) or cannabis and withdrawal from central nervous system depressants (e.g., alcohol and barbiturates) can precipitate a panic attack. However, if panic attacks continue to occur outside of the context of substance use (e.g., long after the effects of intoxication or withdrawal have ended), a diagnosis of panic disorder should be considered. In addition, because panic disorder may precede substance use in some individuals and may be associated with increased substance use especially for purposes of self-medication, a detailed history should be taken to determine if the individual had panic attacks prior to excessive

substance use. If this is the case, a diagnosis of panic disorder should be considered in addition to a diagnosis of substance dependence or abuse. Features such as onset of panic after the age of 45 or the presence of atypical symptoms during a panic attack (e.g., vertigo, loss of consciousness, loss of bladder or bowel control, slurred speech, or amnesia) suggest that a general medical condition or a substance may be causing the panic attack symptoms.

Panic disorder must be distinguished from other mental disorders (e.g., other anxiety disorders and psychotic disorders) that have panic attacks as an associated feature. By definition, panic disorder is characterized by recurrent, unexpected panic attacks, either initially or later in the course, although expected panic attacks may occur as well. In contrast, panic attacks that occur in the context of other anxiety disorders are always expected (e.g., triggered by social situations in social anxiety disorder, by phobic objects or situations in specific phobia, by worry in generalized anxiety disorder, by obsessions in obsessive-compulsive disorder, or by reminders of the trauma in posttraumatic stress disorder).

The focus of the anxiety also helps to differentiate panic disorder from other disorders associated with panic attacks. In panic disorder, the anxiety is about having more panic attacks or their consequences. In other disorders, the panic attacks are symptoms of anxiety about other issues, such as social evaluation in social anxiety disorder, obsessions in obsessive-compulsive disorder, or traumatic experiences in posttraumatic stress disorder.

Comorbidity

The literature is equivocal as to whether depression has an adverse effect on treatment outcome for panic disorder. Several studies have shown that depression at baseline results in poorer outcomes from CBT for panic disorder/agoraphobia (e.g., Rief *et al.*, 2000; Tsao, Lewin, and Craske, 1998). In contrast, other studies have found that depression does not influence response to CBT for panic disorder (e.g., Allen *et al.*, 2010). As such, questions remain regarding how to manage comorbid depression. Some studies (e.g., Craske *et al.*, 2007) would suggest that no adaptation is necessary because targeted CBT will be equally effective with or without the presence of other disorders. Furthermore, according to these studies, comorbid conditions actually improve following targeted panic disorder treatment. On the other hand, these studies typically exclude the most severe depression or suicidal depression. Others would suggest that depression needs to be addressed, such as through the addition of medication, although empirical support for this supposition is lacking.

To date, few studies include Axis II diagnoses in their diagnostic screening procedures perhaps due to the limited availability of reliable and valid methods of assessment. Of those that do, some suggest that individuals with Axis II features have a poorer response to CBT (Hoffart and Hedley, 1997; Marchand *et al.*, 1998). Other studies show the same rate of improvement, although those with Axis II features start and end at a higher level of severity than those without Axis II features (e.g., van den Hout, Brouwers, and Oomen, 2006). Although there are mixed findings regarding the effect of comorbid personality disorders on treatment outcome, it is necessary

for clinicians to be aware of potential adjustments necessary when working with these clients. For instance, clients may exhibit decreased motivation, especially with regards to exposure and between-session homework. Treatment length may need to be increased to allow such clients to reach the same end-state functioning as their non-comorbid counterparts.

Medical comorbidity, such as cardiac arrhythmias or asthma, may slow improvement rates, given the additional complications involved in discriminating anxiety symptomatology from disease symptomatology, increases in actual medical risk, and the stress of physical diseases. However, the effect of medical comorbidity upon CBT outcome has not been assessed to date, and attempts to modify CBT to address medical comorbidity are only just emerging (Lehrer *et al.*, 2008). When working with medical comorbidity, it is recommended that clinicians work closely with physicians and inquire regarding potential contraindications of exposure to interoceptive and external cues given the specific medical condition. Medical “clearance” prior to the onset of exposure therapy is a good standard of practice but is especially important in working with medical comorbidity. Furthermore, in the context of CBT, it is important to help clients learn to articulate the differences between medical symptoms and nonmedical symptoms, and possibly utilize ongoing monitoring of relevant health indices (e.g., blood pressures or heart rate monitors) during exposure therapy. Cognitively, it is important to engage in problem solving around real medical issues.

With respect to substance use disorders, the majority of research studies exclude individuals with comorbid substance disorder. However, one series of case studies (Lehman, Brown, and Barlow, 1998) found that CBT for panic led to reduced alcohol consumption. In the absence of clear research-based guidelines, it is suggested that clinicians strive to understand which disorder is primary and treat that first; so that if substance disorder is primary, it would be treated first while delaying treatment of the panic disorder (Marshall, 1997). Further research on the effects of comorbidity on treatment is desperately needed given the common co-occurrence of anxiety and substance disorders (e.g., Compton *et al.*, 2007).

Efficacy

An extensive body of research has evaluated the efficacy of CBT for panic disorder with or without agoraphobia (PDA). Although agoraphobic avoidance is sometimes associated with less positive response (e.g., Dow *et al.*, 2007), the overall within-group effect size for change in PDA from pre- to posttreatment is very large, for example, $ES = 1.53$ (Norton and Price, 2007). Moreover, the between-group effect size is substantial in comparison to wait-list conditions, for example, $ES = 0.64$ (Haby *et al.*, 2006). However, more research is needed comparing CBT to alternative active treatment conditions.

The effectiveness extends to clients who experience nocturnal panic attacks (Craske *et al.*, 2005b). Furthermore, CBT results in improvements in rates of comorbid anxiety and mood disorders (e.g., Craske *et al.*, 2007; Tsao *et al.*, 2005), although one study suggests that the benefits for comorbid conditions may lessen over time, when assessed 2 years later (Brown, Antony, and Barlow, 1995). Finally, applications of CBT lower

relapse rates upon discontinuation of high-potency benzodiazepines (e.g., Spiegel *et al.*, 1994).

CBT effects generally are sustained over time as meta-analyses show little change (i.e., maintenance of treatment effects) from posttreatment to follow-up, for example, $ES = 0.12$ (Norton and Price, 2007). From their review of meta-analyses for CBT across all disorders, Butler *et al.* (2006) concluded that evidence for maintenance of treatment gains was particularly strong for panic disorder, where the rate of relapse was almost half the rate of relapse following pharmacotherapy. Continuing improvement after acute treatment is facilitated by involvement of significant others in every aspect of treatment for agoraphobia (e.g., Cerny *et al.*, 1987). Also, booster sessions enhance long-term outcomes (Craske *et al.*, 2006).

Efficacy data from research settings are now being complemented by effectiveness data from real-world primary care settings. In a randomized controlled trial in primary care settings with novice therapists, CBT combined with expert recommendations for medication regimens was more effective than treatment as usual (Roy-Byrne *et al.*, 2005). The effects appeared primarily due to CBT (Craske *et al.*, 2005a). In the more recent CALM study (Craske *et al.*, 2011), the effectiveness of CBT for panic disorder in primary care settings was demonstrated in the hands of nonexperienced therapists with the aid of a computerized guide, combined with expert recommendations for medication, relative to treatment as usual.

Even though CBT for panic disorder/agoraphobia is efficacious and effective, there is room for improvement. One study estimated that 30% of clients continued to function poorly at follow-up and only 48% reached high end-state status (Brown and Barlow, 1995). In a landmark study (Barlow *et al.*, 2000), only 32% of panic disorder clients assigned to CBT alone demonstrated strong treatment response 12 months after acute treatment. Finally, of those who do start treatment, the mean dropout rate from CBT for panic disorder/agoraphobia is 19%, with a range of 0–54% (Haby *et al.*, 2006).

Medication Management

The combination of CBT (or in some cases behavioral therapy) and antidepressant treatment has shown a small advantage over either CBT alone or antidepressants alone for panic disorder/agoraphobia. A meta-analysis of 21 trials with 1709 clients compared either CBT or antidepressant treatment for panic disorder/agoraphobia to combined CBT-antidepressant treatment (Laberge *et al.*, 1993). Combined treatment was superior to antidepressant pharmacotherapy (RR 1.24, 95% CI 1.02–1.52) or psychotherapy (RR 1.17, 95% CI 1.05–1.31). However, combined treatment resulted in a somewhat greater number of client dropouts due to side effects than psychotherapy alone. The superiority of the combination over either monotherapy appeared to persist as long as the drug was continued. Following discontinuation of the antidepressant, the groups initially assigned to either combined treatment or CBT did better than clients assigned to pharmacotherapy. Similarly, in the landmark study mentioned earlier (Barlow *et al.*, 2000), following medication discontinuation, the combination of medication and CBT fared worse than CBT alone,

suggesting the possibility that state or context-dependent learning in the presence of the medication may have attenuated the new learning that occurs during CBT. For this reason, it is generally recommended that CBT booster sessions be conducted during and following medication withdrawal, although direct empirical testing has not been conducted.

Findings from the combination of high-potency benzodiazepines with behavioral treatments for agoraphobia are limited and mixed (Marks *et al.*, 1993; Wardle *et al.*, 1994). However, several studies reliably show that chronic use of high-potency benzodiazepines for panic disorder/agoraphobia, initiated well before CBT, has detrimental effects on short- and long-term outcomes from CBT, including more attrition, less improvement, and greater likelihood of relapse (e.g., Westra, Stewart, and Conrad, 2002).

Adaptations for Ethnic Subgroups

In general, CBT is aligned with European and North American values of change, self-disclosure, independence and autonomy, and rational thinking – all of which are at odds with values of harmony, family and collectivism, and spirituality that define many other cultures (see Hays and Iwamasa, 2006). Also, cognitive strategies are closely aligned with the European/North American value of rational thinking. As noted by Hays and Iwamasa (2006), emphasis upon cognition, logic, verbal skills and rational thinking can undercut the value of spirituality in many cultures. Related is the emphasis of cognitive strategies upon reductionist cause and effect relations. In contrast, Asian cultural beliefs, for example, emphasize balance (or yin and yang), evaluation of systems holistically, and indirect causes for events. Thus, a client from a different culture may take into account additional causes for events that are not immediately obvious to a North American/European therapist, thereby potentially undercutting the latter's attempts at cognitive restructuring. Another issue is the locus of control, which tends to be more internal than external in Caucasians and more external than internal in many Asian cultures, wherein certain events are judged to be unrelated to one's own actions. In addition, whereas a typical target of CBT is negative self-statements, such self-criticism may be viewed as a motivator to achieve in other cultures, such as Chinese cultures (Hwang and Wood, 2007). Thus, what a therapist may judge to be a "logical outcome" may not coincide with the client's beliefs. Cultural sensitivity of cognitive strategies mandates that therapists become knowledgeable about the client's cultural values and beliefs, something that would be achieved through functional analyses.

To adapt CBT to different cultural groups may mean that exposure therapy goals of independence and autonomy (e.g., being able to drive alone and travel far from home) are cast within, and not at conflict with, cultural values that emphasize family and collectivism. Furthermore, cognitive therapy may be modified for cultures that emphasize balance and indirect causes of events rather than reductionist cause-effect relations. That is, the method by which cognitive restructuring is most effectively implemented may differ across cultures, with Western culture being more suited to single overriding alternative interpretation of a given event and Eastern culture being