

Lothar Staeck

# Fascination Amazon River

Its People, Its Animals, Its Plants

 Springer

# Fascination Amazon River



Geographical overview of the fascinating world of the Amazon river from Nauta in Peru to the estuary

Lothar Staeck

# Fascination Amazon River

Its People, Its Animals, Its Plants

 Springer

Lothar Staeck  
Berlin, Germany

The translation was done with the help of artificial intelligence (machine translation by the service DeepL.com). A subsequent human revision was done primarily in terms of content.

ISBN 978-3-662-64451-5                      ISBN 978-3-662-64452-2 (eBook)  
<https://doi.org/10.1007/978-3-662-64452-2>

© Springer-Verlag GmbH Germany, part of Springer Nature 2022

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, expressed or implied, with respect to the material contained herein or for any errors or omissions that may have been made. The publisher remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This Springer imprint is published by the registered company Springer-Verlag GmbH, DE part of Springer Nature.

The registered company address is: Heidelberger Platz 3, 14197 Berlin, Germany

# Contents

<b>1</b>	<b>Introduction: What Travellers Can Really See on Their Trip to the Rainforest</b>	<b>1</b>
<b>2</b>	<b>The Amazon: The Largest River on Earth</b>	<b>5</b>
2.1	The River Between Its Source and Mouth	7
2.2	Várzeas: Floodplain Meadows	12
2.3	Igapós: Floodplain Forests	14
2.4	River Meanders and Terra Firme	16
2.5	White Water	22
2.6	Black Water	23
2.7	Clear Water	24
<b>3</b>	<b>Cities Along the Amazon River</b>	<b>29</b>
3.1	Iquitos	29
3.2	Leticia	32
3.3	Manaus	34
3.4	Parintins	39
3.5	Santarém	40
3.6	Belém	41
<b>4</b>	<b>The People of the Amazon River</b>	<b>45</b>
4.1	The Original Population	45
4.2	The caboclos	68

<b>5</b>	<b>The Rainforest</b>	77
5.1	Greatest Diversity of Plant Species	77
5.2	A “New” Primary Forest Cannot Be Planted	79
5.3	The Ground Layer of the Rainforest	80
5.4	Middle Floor of the Rainforest	86
5.5	The Soil of the Old-Growth Forest	88
5.6	Top Floor of the Rainforest	92
5.7	Root Types	94
5.8	The Brazil Nut Tree	96
5.9	Epiphytes and Climbers	100
5.10	Summary: Structure of the Rainforest	103
<b>6</b>	<b>Flowering Plants in and Around Water</b>	107
6.1	Trees	108
6.2	Shrubs	127
6.3	Perennials	136
6.4	Aquatic Plants	141
6.5	Epiphytes	159
6.6	Vines	167
<b>7</b>	<b>Flowering Plants in the Rainforest</b>	171
7.1	Introduction	171
7.2	Deciduous Trees of the Terra Firme	172
7.3	Palms	189
7.4	Giant Strelitzia (Strelitziaceae)	193
<b>8</b>	<b>Animal Observations by and in the Water</b>	195
8.1	Introduction	195
8.2	Fish	196
8.3	Frogs	225
8.4	Reptiles	230
8.5	Birds	242
8.6	Mammals	279
8.7	Butterflies	290
<b>9</b>	<b>Animal Observations in the Rainforest</b>	297
9.1	Poison Dart Frogs	298
9.2	Lizards	299
9.3	Parrots	301
9.4	Toucans	305

9.5	Woodpeckers	306
9.6	Tanagers	306
9.7	Monkeys	309
9.8	Porcupine Relatives	312
9.9	Tarantulas	314
9.10	Ants	316
9.11	Termites	321
9.12	Beetles and Praying Mantises	323
<b>10</b>	<b>Concluding Note: What Is the Future of the Amazon Lowlands?</b>	
		327
10.1	Forest Destruction	327
10.2	Climate Change	328
10.3	Population Growth	329
	<b>Index</b>	333



# List of Figures

Fig. 2.1	Size comparison of Amazonia and the land area of the United States (without Alaska and Hawaii)—Amazon basin is marked in green colour	6
Fig. 2.2	Course of the primeval Amazon about 130 million years ago	8
Fig. 2.3	Remains of a laterite cliff of the proto-Amazon with breeding burrows of the <i>Ringed kingfisher</i> (Sect. 8.5.6)	10
Fig. 2.4	Grass-reed islands drifting downstream during the rainy season	11
Fig. 2.5	<i>Várzea</i> lakescape of floodplain meadows; the arrow marks the border of the riverbed	13
Fig. 2.6	Huge masses of water flow into the flat <i>hinterland</i> , where they form the <i>várzeas</i> . In the background on the right, numerous giant water lilies ( <i>Victoria amazonica</i> ) grow	14
Fig. 2.7	The flooded forests of the Blackwater rivers and lakes (Blackwater Igapós) are under water for up to 7/8 months	15
Fig. 2.8	Schematic representation of river cliffs and slip-off slopes: as the rainy season progresses, the river cliffs are eroded and bank failures occur. During the dry season, sediments are deposited on the slip-off slopes, so that sandbanks can gradually form	16
Fig. 2.9	In the “death zone” of the river cliff, many trees fall into the river every year	17
Fig. 2.10	Schematic representation of the water breakthrough at a river bend	18
Fig. 2.11	During the dry season, a steep bank rises up to 15 m high at the river edge	19
Fig. 2.12	Bank edge at low water	19
Fig. 2.13	Riparian edge during floods	20
Fig. 2.14	Cross-section of the river landscape with its ecosystems on both sides of the Amazon River	20
Fig. 2.15	Cattle transport on the Amazon River	21

Fig. 2.16	Typical multi-storey Amazon ship	22
Fig. 2.17	Distribution of the three types of water bodies in the Amazon basin: 1 = Ancient degraded uplands (Guyanas and central Brazil), 2 = Swampy lowlands, floodplains and pale sands, 3 = Floodplain forests (white-water Igapós) and terra-firme on alluvial sediments of the Andes, 4 = Rainforests (terra-firme), 5 = Floodplain forests (mostly black-water Igapós)	23
Fig. 2.18	With its typical clay-yellow coloration, the Amazon is a white-water river with its typical clay-yellow coloration	24
Fig. 2.19	(a) A typical blackwater river with the characteristic cola-like colouring, the tannins sometimes staining the substrate reddish. (b) The <i>Rio Tapajos</i> , a typical clear-water river, which appears light green to yellow-green due to algae growth on the substrate	26
Fig. 2.20	Schematic representation of acidification in lakes near the Amazon	27
Fig. 3.1	More than 40,000 three-wheeled <i>motokars</i> drive through the streets of Iquitos	30
Fig. 3.2	The fat <i>suri</i> grubs of the palm weevil are considered a delicacy by the natives of Iquitos	31
Fig. 3.3	Boundary between <i>Leticia</i> (Colombia), <i>Tabatinga</i> (Brazil) and <i>Santa Rosa</i> (Peru)	32
Fig. 3.4	The main street <i>Calle 8</i> in <i>Leticia</i>	33
Fig. 3.5	Confluence of the <i>Rio Solimões</i> (white-water river) and the <i>Rio Negro</i> (black-water river) near <i>Manaus</i> . From here on the river is called “Amazonas” by the Brazilians	35
Fig. 3.6	The now renovated Fish Hall, built in 1883 according to the model of <i>Les Halles</i> in Paris, was designed by Gustave Eiffel	36
Fig. 3.7	The <i>Manaus</i> Opera House was built in 1896 at the height of the rubber boom	38
Fig. 3.8	In the dry season there are snow-white sandy beaches in <i>Alter do Chão</i>	41
Fig. 3.9	The <i>Ver-o-peso</i> market in <i>Belém</i>	42
Fig. 4.1	A member of the <i>Ticuna</i> Indians in traditional dress	46
Fig. 4.2	Members of the same ethnic group—dressed in modern clothes, they belong to the clan of unwinged animals, namely the jaguar clan	47
Fig. 4.3	The indigenous population of the Amazon shows great genetic similarities with people living today in New Guinea, Australia and the Andaman Islands—the darker the colour circles, the closer the relationship	47
Fig. 4.4	<i>Terra Preta</i> discoveries in the middle and lower Amazon	48
Fig. 4.5	At the end of the sixteenth century, pictures arrived in Europe that were supposed to show members of the <i>Ewaipanoma</i> Indians from Amazonia: headless beings with facial drawings on their chests	49

Fig. 4.6	The indigenous people believed in animate nature. At ritualized festivals, Indians dress up with demon and animal masks	50
Fig. 4.7	<i>Yagua</i> women in their typical traditional clothing	51
Fig. 4.8	Boys of the <i>Yagua</i> tribe present their “pets”: a parrotlet, a small squirrel monkey and a small spectacled caiman	52
Fig. 4.9	The clothing of the <i>Huitotos</i> is traditionally made from beaten bark bast of a fig species. The upper parts of the body are dyed black with the sap of the genipapo tree	53
Fig. 4.10	(a) Huitoto woman today, (b) Huitoto woman around 1870	53
Fig. 4.11	After the bath the <i>Ticuna</i> girl is decorated with <i>genipapo</i> juice and bird feathers	54
Fig. 4.12	The girls are led into the meeting house ...	55
Fig. 4.13	... where female relatives pull out their head hair in clumps	56
Fig. 4.14	<i>Ticuna</i> juvenile with filed teeth resembling piranha dentition	57
Fig. 4.15	Dance performance by <i>Desano Indian</i> men in traditional dress, with necklace of caiman canine teeth	58
Fig. 4.16	Young <i>Desano</i> woman with typical body painting	59
Fig. 4.17	<i>Desano Indian</i> with feather crown made of macaw feathers	60
Fig. 4.18	(a) Hut of the indigenous people in traditional construction, (b) <i>Maloca</i> (community house) of the <i>Yagua</i> and <i>Bora Indians</i>	61
Fig. 4.19	Principle of <i>shifting cultivation</i> : cleared piece of rainforest, tree trunks and roots left lying around	62
Fig. 4.20	(a) Intact forest with a closed canopy cover, (b) Trees are completely removed by caboclos and white settlers, including trunks and roots, (c) Massive soil erosion takes place after only a short time	63
Fig. 4.21	Wood of the muirapiranga tree, also called bloodwood	65
Fig. 4.22	(a) Fruits of the genipapo tree, (b) Prepared and oxidized juice of genipapo fruits	66
Fig. 4.23	Inflorescence of the piaõ bush	67
Fig. 4.24	The berries of the <i>camu camu</i> bush have 40 times the vitamin C content of oranges	67
Fig. 4.25	The nux devil doer ( <i>Strychnos toxifera</i> ) contains the alkaloid toxiferin in its seeds, which is highly toxic	69
Fig. 4.26	(a) <i>Caboclos</i> are descendants of Indians and European immigrants, (b) The faces of adult <i>caboclos</i> are brown, the hair colour is consistently black	70
Fig. 4.27	A <i>caboclo</i> hut surrounded by a palm plantation (açai palms)	70
Fig. 4.28	In cassava, the spindle-shaped tubers develop from the lateral roots. In the background, the hand-shaped leaves can be seen on the shoots	71
Fig. 4.29	(a) After peeling, the cassava tuber must be soaked for 2 days, during which the prussic acid contained in the tubers is released	

	into the water, <b>(b)</b> After soaking, the soft cassava tubers are pressed through a basket weaving ( <i>tipiti</i> )	72
Fig. 4.30	The moist cassava mass is roasted on the pan to make flour	72
Fig. 4.31	The slender açai palms, which grow to a height of 20 m, only grow in the Amazon basin	73
Fig. 4.32	<b>(a)</b> The fruit stand of the açai palm with fruits that resemble blueberries in size and colour, <b>(b)</b> Açai juice	74
Fig. 5.1	Dwarfed remaining tree saplings (referred to by me as “oskars”) at the bottom of the primary forest	81
Fig. 5.2	<b>(a)</b> The red-yellow flower colour of the sore-mouth-bush ( <i>Psychotria poeppigiana</i> ) signals a nectar source to butterflies and birds, <b>(b)</b> The blue fruits of the same plant attract birds in the dark jungle	82
Fig. 5.3	The bright red sepals of these plants have the nickname “Lábias da prostituta” in Brazil	83
Fig. 5.4	Mushrooms ( <i>Basidiomycetes</i> ) with colourful fruiting bodies recycle leaves ..	83
Fig. 5.5	... and rotten wood; this picture shows a group of the species <i>Cookeina speciosa</i>	84
Fig. 5.6	<b>(a)</b> The leaves of a begonia ( <i>Begonia nasoniana</i> ) pretend to be old and eaten away, <b>(b)</b> In representatives of the <i>Calathea</i> and <i>Maranta</i> species (here <i>Calathea elliptica</i> ), the leaf contours are blurred by coloured sprinkles to deceive leaf-eating insects	85
Fig. 5.7	The river bank, densely overgrown with pioneer woodland, gives the (false) impression that the primeval forest is impenetrable	86
Fig. 5.8	<i>Cauliflory</i> is quite widespread in the rainforest: this picture shows wild cacao ( <i>Herrania mariaae</i> )	87
Fig. 5.9	Fresh young leaves hanging limply to fool leaf-eating insects	88
Fig. 5.10	For a perfect illusion even false colours (here purple) are used	89
Fig. 5.11	<b>(a)</b> Under the thin humus layer extends the climax soil (latsole), which is forever infertile, <b>(b)</b> This ochre-red hard soil structure is called laterite. It is also forever infertile	90
Fig. 5.12	Subterranean symbiotic network between the roots of rainforest trees and <i>mycorrhizae</i> (fungal filament network). Large star: old rainforest giant; small stars: younger trees	91
Fig. 5.13	The giant trees (here a kapok tree), which can reach heights of up to 60 m, tower above the closed canopy	93
Fig. 5.14	The jungle giants develop board roots up to 9 m high, which stabilise the tree like a foundation	94
Fig. 5.15	<i>Socratea</i> palms (“running palms”) develop tall stilt roots that increase their stability and enable the tree to move even in the direction of the arrow	95
Fig. 5.16	Shoot-born roots originating from the base of the stem, as in the case of açai palm	96

Fig. 5.17	The slender Brazil nut tree grows up to 50 m high	97
Fig. 5.18	The flowers of the Brazil nut tree	97
Fig. 5.19	(a) Seedlings grow from the capsule fruit of the Brazil nut tree and compete with each other. Only one will survive. (b) Inside the almost spherical fruit of the Brazil nut lie up to 32 hard-shelled tasty seeds. However, the woody seed coat still has to be removed	98
Fig. 5.20	Different <i>epiphyte</i> —species on a host tree. In the foreground, flowering representatives of the <i>Bromeliaceae</i> (pineapple family) can be seen	100
Fig. 5.21	In an old liana shoot, thin strands of tissue, often separated from each other, are united—like bundles of cables with thin wires, so they are flexible and tough	102
Fig. 5.22	A strangler fig at the beginning of its growth: tentacle-like, it begins to grow around its host	103
Fig. 5.23	A strangler fig has already almost completely enclosed its host tree	104
Fig. 5.24	After many decades, the battle is over: the strangler fig has taken the place of its host. The hollow cylinder marks the space of the former host tree	105
Fig. 5.25	Structure of the Amazon rainforest	105
Fig. 6.1	The shield-leaved snake wood is a pioneer plant with a high light requirement	109
Fig. 6.2	The <i>Gustavia tree</i> has delicately fragrant, splendid flowers	111
Fig. 6.3	The attractive flowers of the <i>Eschweilera tree</i> are captivating because of their numerous, ring-shaped, lemon-yellow stamens, which are located in the centre of the cream-coloured petals	112
Fig. 6.4	With the onset of the rainy season, innumerable 25 cm long woody pods hang from the <i>Campsiandra tree</i>	113
Fig. 6.5	The flowers of the <i>Campsiandra tree</i> , which are blood-red before flowering, are densely packed together in an inflorescence	114
Fig. 6.6	The mushroom-like inflorescence of the <i>Parkia tree</i> is unique	115
Fig. 6.7	The fruits of the <i>Parkia tree</i> grow into 15 cm long, red-brown pods	116
Fig. 6.8	The decorative flowers of the Guiana chestnut resemble a shaving brush. Its fruits, up to 30 cm long, harbour numerous edible nuts	117
Fig. 6.9	The imposing reddish-brown fruits of the Munguba tree still hang on the tree when the leaves have already fallen off	118
Fig. 6.10	The creamy white flowers of the Munguba tree seem to consist only of stamens	118
Fig. 6.11	Because of the shape of its flowers, this tree is called the “shameless” or <i>clitoria</i>	120
Fig. 6.12	(a) The filigree flowers of the <i>Inga tree</i> , (b) The slender fruits, up to 50 cm long, are the reason for the popular name “monkey’s tail”	120

**xiv**      **List of Figures**

Fig. 6.13	The Marimari tree has large yellow flowers with conspicuous stamens	122
Fig. 6.14	The green tasty pulp surrounds the ring-shaped seeds of the Marimari tree	122
Fig. 6.15	(a) Inflorescence of the garlic pear, (b) The round, slightly garlic-scented fruits of the garlic pear are readily eaten by fish	123
Fig. 6.16	The striking, large, red-brown fruit clusters hang from the mighty Morich palm. The edible, decorative fruits are covered with large scales, which can be peeled off with the fingernail after watering	125
Fig. 6.17	The buçu palm has huge leaflets up to 10 m long	127
Fig. 6.18	The two black fruits of the Mickey Mouse bush look like the ears of the comic strip character of the same name. The additional drawing on the red flower base completes the impression of a Mickey Mouse figure	128
Fig. 6.19	The flowers of the <i>Heisteria bush</i> look similar to the Mickey Mouse bush	129
Fig. 6.20	(a) The wild form of the guaraná shrub <i>Paullinia pinnata</i> , (b) The fruits of the “real” guaraná look like eyes	130
Fig. 6.21	(a) The golden trumpet has large flowers resembling the shape of the brass instrument, (b) The sister of the golden trumpet, <i>Allamanda violacea</i> , grows as a delicate-limbed climber in floodplain meadows	132
Fig. 6.22	(a) Blue potato shrub <i>Solanum crinitum</i> has curled petals, (b) Because of its bizarre udder-shaped form, the species <i>Solanum mammosum</i> is often planted in gardens	133
Fig. 6.23	Because of the numerous, upright stamens, the Amazon residents call this plant “monkey’s brush”. In the related species, <i>Combretum cacoucia</i> , the flowers hang in clusters on the branch	134
Fig. 6.24	The almost palm-sized flowers of <i>Odontadenia hypoglauca</i> have a colour pattern in their centre that looks like a second flower	136
Fig. 6.25	The large red-yellow bracts of the hanging lobster claw hang down like garlands	137
Fig. 6.26	The mighty pseudo-stems of the arum lily push out of the bank mud like asparagus tips	138
Fig. 6.27	The inflorescence of the arum lily consists of white bracts and a fleshy, cream-coloured flower bulb. In order to attract the beetles required for pollination, the temperature of the flower bulb increases to over 40 degrees Celsius (see the red colour in the picture!)	139
Fig. 6.28	The flowers of the Dormilones are pink to light purple	141
Fig. 6.29	Up to 20 different aquatic plants grow in the floodplain meadows	142

- Fig. 6.30 In the calm and nutrient-rich floodplain meadows, Queen Victoria's water lily can develop magnificently 143
- Fig. 6.31 The underside of the leaves of Queen Victoria's water lily is armed with centimetre-long spines to keep predators away 144
- Fig. 6.32 At dusk on day 1, the Queen Victoria's water lily blooms snow-white 145
- Fig. 6.33 Different beetle species are attracted by the scent, high temperatures and the candle-light colour of the flower and crawl into the flower interior 145
- Fig. 6.34 At the end of day 2, the flower has changed to a dark red-purple color 146
- Fig. 6.35 Characteristic of the common water hyacinth is the bright yellow spot on the upper floral lobes 148
- Fig. 6.36 Round-leaved tropical pickerelweed (*Pontederia rotundifolia*) flowers in a spikey, pale pink inflorescence only 4 cm across 149
- Fig. 6.37 The light green leaf rosette is characteristic of water cabbage. The tiny inflorescence consists of a white bract and an equally tiny yellowish flower bulb 150
- Fig. 6.38 (a) The leaves of the floating heart look like a mini water lily, but its decorative flowers appear much more delicate and smaller, (b) The mosaic flower forms pretty geometric patterns on the water surface, (c) The clearly visible white breathing roots next to the flower of the floating Ludwigia give this plant a fantastic buoyancy. A few water ferns (*Carolina azolla*) float next to the flower 152
- Fig. 6.39 The leafy bladderwort is a carnivorous plant. It catches its prey, e.g. water fleas, with numerous catch bladders 154
- Fig. 6.40 The eared moss consists of only two ear-shaped floating leaves. On the upper side of the leaves there are countless whisk-shaped papillae, which keep the leaves on the water surface at all times (1  $\mu\text{m}$  = 0.001 mm) 156
- Fig. 6.41 The *Carolina azolla fern* is only 5–10 mm in size 158
- Fig. 6.42 The ruby-lipped *Cattleya* is the national flower of Brazil 160
- Fig. 6.43 The parrot flower is pollinated by hummingbirds 161
- Fig. 6.44 (a) This tillandsia (*Tillandsia paraensis*) likes to grow as an epiphyte on dead trees on the banks of blackwater rivers, (b) This tillandsia (*Tillandsia limbata*) forms a bizarre branched inflorescence, (c) This lance rosette lives epiphytically and also terrestrially—i.e. “rooting in the ground”, (d) This *Guzmania* (*Guzmania sanguinea*) lives epiphytically on its carrier tree 163
- Fig. 6.45 The large philodendron lives initially as an epiphyte developing numerous aerial roots. The cream-coloured flower bulb is surrounded by a decorative bract. The snakelike curved fruit of the

	philodendron is densely packed with reddish-purple, fragrant berries that attract many species of birds	165
Fig. 6.46	The heartleaf vine is a climber with decorative yellow, orchid-like flowers	167
Fig. 6.47	(a) The red passion flower likes to grow on sunny riverbanks ( <i>Passiflora cochinea</i> ), (b) The passionflower <i>Passiflora vitifolia</i> has vine-like stem leaves and a secondary crown of white-purple appendages, (c) Passionflower <i>Passiflora laurifolia</i> has laurel-like stem leaves and a purple-white banded subcrown, (d) Passionflower <i>Passiflora nitida</i> has a long, deep blue subcrown with white curled filaments in the last third	169
Fig. 6.48	The male flower of the jungle cucumber is densely covered with silky hairs	170
Fig. 7.1	The yellow trumpet tree flowers in the dry season when the deciduous leaves are shed	173
Fig. 7.2	The funnel-shaped flowers of the pink trumpet tree	173
Fig. 7.3	The fruits of the cannonball tree, which are the size of a child's head, are one of the favourite foods of wild boars	175
Fig. 7.4	The bizarre flowers of the cannonball tree are cauliflorous, i.e. they grow directly out of the trunk	176
Fig. 7.5	Like all flowers of the <i>Lecythidaceae</i> , the flowers of the monkey pot are also peculiarly shaped. The up to 30 cm large fruits of this tree contain numerous tasty seeds	177
Fig. 7.6	The cauliflorous flowers of the cacao tree are tiny, but the fruits are surprisingly large at 15 cm. They have a leathery-hard dark yellow to reddish colour	178
Fig. 7.7	The fruit of the <i>cupuaçu tree</i> , a relative of the cacao tree, contains an aromatic pulp	180
Fig. 7.8	The seeds of the lipstick tree contain a scarlet dye that plays an important role in the culture of the Yaguas	182
Fig. 7.9	The unusual inflorescence of the tree "Pride of Trinidad and Tabago" consists of numerous bright red bracts up to 10 cm long, which stand on an axis up to 80 cm long. Between them sit the inconspicuous flowers	183
Fig. 7.10	Numerous aerial roots hanging down from the main stem as well as thick secondary stems already anchored in the soil are characteristic of the Florida strangler fig. The black colouring of the stems marks the high water line	184
Fig. 7.11	The rubber tree has brown speckled seeds	186
Fig. 7.12	The rubber tree's gummy milky sap flows out of the bark from a diagonally applied cut	187
Fig. 7.13	The golden shower tree of the Amazon basin confirms Alfred Wegener's theory of continental drift	190



Fig. 7.14	The fruits of the peach palm provide a popular starchy vegetable	191
Fig. 7.15	The oily flesh of the tucumã palm, which is only a few millimeters thick, has an aromatic, nutty flavor	192
Fig. 7.16	The huge greenish-yellow inflorescence of the giant strelitzia resembles the natal strelitzias from South Africa	193
Fig. 8.1	The peacock bass is a voracious predator	199
Fig. 8.2	In the marble cichlid, the bright orange-red spots and stripes on the sides of the body and the eye dummies on the rear fin rays are particularly striking	200
Fig. 8.3	The Heckel's cichlid is a popular aquarium fish	202
Fig. 8.4	(a) The leopard pike cichlid is a voracious prey snapper, (b) At vegetated bank sections or in the dry season, numerous fish species can be caught in a short time with towed nets	203
Fig. 8.5	The most famous fish in the Amazon is the red piranha	205
Fig. 8.6	The black piranha weighs 2 kg and bites with 30 times its own weight	206
Fig. 8.7	The pirapitinga is called "false fish" by residents because, as a juvenile, it is strikingly similar in colour and shape to the red piranha	208
Fig. 8.8	The black pacú is an impressive fish, with up to 1.10 m in body length and weighing 40 kg. Its teeth sit in the lower and upper jaw in two rows behind each other	209
Fig. 8.9	The flag-tailed prochilodus is an important food fish in the <i>Manaus</i> region	210
Fig. 8.10	The "cachorro" (Brazilian for "dog") has a strong dentition of conically shaped teeth	212
Fig. 8.11	Sailfin catfish are armoured all over with bone shields	213
Fig. 8.12	The redtailed catfish is one of the impressive fishes of white water, with a body length of up to 1.30 m	214
Fig. 8.13	The tiger-like colour pattern of the tiger surubim is characteristic of this popular, boneless food fish	215
Fig. 8.14	With its flank spines and scalpel-like ventral fin spines, the rip saw catfish (Cuiú-Cuiú) is even safe from jaguars and caimans	216
Fig. 8.15	Candirús invade the intestines of their host, where they feed on their blood. It also happens that they enter the body orifices of humans	217
Fig. 8.16	The silver arowana ("water monkey") jumps up to 2 m out of the water to snatch its prey on branches near the shore	218
Fig. 8.17	The whip-like tail of this freshwater stingray cururu has barbed spines. In case of danger, they release a life-threatening poison via these spines. The small picture shows an incompletely developed stingray with a large yolk sac	220

Fig. 8.18	The electric eel emits short bursts of more than 600 V to catch prey and to defend itself. Its “lungs” with numerous well perfused papillae are located on the roof of its palate	221
Fig. 8.19	One of the largest freshwater fish in the world is the Arapaima (Pirarucu), with a body length of more than 3 m	222
Fig. 8.20	This giant pirarucu (= red fish) was photographed 30 years ago at the <i>Manaus</i> fish market	224
Fig. 8.21	Frogs are masters of camouflage. Here The tree frog in this photo is hardly recognizable	225
Fig. 8.22	The dotted treefrog is nocturnal, as its large eyes reveal	227
Fig. 8.23	The calls of the blacksmith treefrog sound like hammer blows on an anvil	228
Fig. 8.24	The tarsier leaf frog is perfectly adapted in colour to its environment	229
Fig. 8.25	Bullfrogs inflate their lungs in danger, enlarging their bodies and standing up on their four limbs	230
Fig. 8.26	The green ameiva lizard hunts insects and spiders	232
Fig. 8.27	The male of the green iguana adorns himself with a large dorsal crest	233
Fig. 8.28	A bony transverse bar between the corners of the eyes helped give the spectacled caiman its name. The small illustration shows a young spectacled caiman	234
Fig. 8.29	(a) With a body length of up to 9 m, the anaconda is the largest snake in the world, (b) With their deeply split tongues, anacondas sniff out even the slightest scent of their prey	237
Fig. 8.30	The <i>boa constrictor</i> is a lurking hunter	239
Fig. 8.31	The rainbow boa is covered with iridescent scales that give rise to rainbow-like reflex patterns	240
Fig. 8.32	The Amazon tree boa is a night hunter with deeply recurved, needle-shaped fangs	241
Fig. 8.33	Pyramidal horny shields on the dorsal carapace of the mata-mata	242
Fig. 8.34	The large-billed tern is ubiquitous in the Amazon river system	243
Fig. 8.35	The wattled jacana lives between the floating leaves of the <i>várzeas</i>	244
Fig. 8.36	(a) In the great egret—here in courtship dress—the bill is yellow, and the legs and feet are black, (b) In the snowy egret it is the other way round: black bill and yellow feet	246
Fig. 8.37	At 1.30 m, the cocoi heron is the largest heron in the Amazon lowlands	247
Fig. 8.38	The capped heron is unmistakable with its long white decorative feathers on the head and blue face	248
Fig. 8.39	The young birds of the rufescent tiger herons have tiger-like feather markings	249
Fig. 8.40	The adult rufescent tiger heron is very conspicuously coloured	250

Fig. 8.41	The mangrove heron places insects as bait on the water surface to attract prey fish	251
Fig. 8.42	The catfish is rotated in its beak until its razor-sharp fin rays cannot injure the olivaceous cormorant when it is swallowed down	252
Fig. 8.43	The darter is a relative of the cormorants	253
Fig. 8.44	The breast of the male ringed kingfisher is reddish-brown, the plumage otherwise rather grey (size: 42 cm)	254
Fig. 8.45	The male Amazon kingfisher has a reddish-brown breast and metallic green plumage (size: 30 cm)	255
Fig. 8.46	The breast of this male green kingfisher is not yet reddish-brown, the plumage is green with small white spots on the wings (Size: 20–25 cm)	256
Fig. 8.47	The osprey has a bright white coloured body underside	256
Fig. 8.48	The roadside hawk is only about 40 cm tall	258
Fig. 8.49	The savannah hawk is easily recognised by its reddish-brown feathering	258
Fig. 8.50	The black-collared hawk is unmistakable by its white head above the cinnamon-coloured body and the black half-ring on the neck	259
Fig. 8.51	The snail kite has distinctive ash-grey or grey-blue plumage. It is specialized on apple snails	260
Fig. 8.52	The yellow-headed caracara is a food opportunist	261
Fig. 8.53	The black caracara has glossy black plumage	262
Fig. 8.54	In black vultures, the short wings prevent elegant, aerodynamic flying	263
Fig. 8.55	The bright red, naked head gives the turkey vulture its name. In contrast, the yellow-headed vulture (small image) has a brightly coloured naked head, with the yellow parts of the face dominating	264
Fig. 8.56	King vultures have a prominent, wart-like bulging skin flap above the beak	265
Fig. 8.57	The hoatzins communicate with each other with a hoarse buzzing sound	266
Fig. 8.58	In the horned screamers a thin, whitish, horn-like structure grows on the forehead and on the wings there are two pointed spines	268
Fig. 8.59	Hummingbirds, such as the plain-bellied emerald must visit up to 3000 flowers a day to avoid starvation	269
Fig. 8.60	The greater ani has a prominent hump on the upper beak	271
Fig. 8.61	In the smooth-billed ani the upper beak is raised like a comb	271
Fig. 8.62	(a) Breeding colony of yellow-rumped caciques, (b) Breeding colony of crested oropendola	272

Fig. 8.63	Yellow-rumped caciques are black in colour except for yellow patches on the wings, rump and tail	273
Fig. 8.64	The crested oropendolas are black in colour except for the maroon root of the tail and the yellow butt	274
Fig. 8.65	Great kiskadees are called <i>Bem-ti-vi</i> (= "I saw you well") in Brazilian	275
Fig. 8.66	The tail of the male fork-tailed flycatcher is three times as long as its body	276
Fig. 8.67	<b>The</b> male of the marsh tyrant has a white head with otherwise black plumage	276
Fig. 8.68	The common potoos are perfectly adapted to their environment—during the day and also at night	277
Fig. 8.69	The common nighthawk flies over the water surface in an irregular zigzag flight	279
Fig. 8.70	The pink river dolphin is called "boto" by the river people	280
Fig. 8.71	The Tucuxi dolphin (or <i>Sotalia</i> ) is similar in appearance to the marine dolphins	282
Fig. 8.72	The fur of sloths is remarkably dense and shaggy	284
Fig. 8.73	The mask-like facial markings with black snout and dark eyes is a distinctive feature of the squirrel monkeys	286
Fig. 8.74	The strong jaws of the white-fronted capuchin give it a fierce appearance	288
Fig. 8.75	The piercing, lion-like call of the howler monkeys serves to mark their territory	289
Fig. 8.76	The upper wing surfaces of the peleides blue morpho are bright blue. This colour impression is created by <i>interference</i> which refracts the structure of the wing scales so that only the blue light components are visible. When this butterfly is sitting, only the dark colouring of its underwings with the seven eye dummies is visible	291
Fig. 8.77	The caterpillar of the bull's eye moth protects itself with innumerable poisonous stinging hairs	292
Fig. 8.78	The caterpillar of the flannel moth is surrounded by a tangle of long, poisonous hairs	293
Fig. 8.79	The bristly hairs of the caterpillar of the wool moth release a nerve toxin when touched. The head is on the left side of the photo	294
Fig. 8.80	The caterpillars of the tetrio phynx wear a spectacular warning costume	294
Fig. 9.1	The belly-spotted poison dart frog signals its venomousness with its striking colour pattern	298
Fig. 9.2	The red-backed poison dart frog, only 12 mm in size, is also brightly coloured	300

Fig. 9.3	(a) The striped basilisk can run up to four metres across the water in case of danger, (b) The Amazonian anolis can change its basic body colouring like a chameleon	301
Fig. 9.4	(a) At 90 cm, the scarlet macaw is the largest parrot in the Amazon. (b) The blue-and-yellow macaw also lives in the Amazon lowlands	303
Fig. 9.5	The blue-fronted Amazon parrot belongs to the Amazon parrots. The tui parakeet (smaller picture) belongs to the true parrots	304
Fig. 9.6	The giant toucan ( <i>toco</i> ) has a 20 cm long, red-orange coloured beak	305
Fig. 9.7	With its 38 cm, the powerful woodpecker is one of the large woodpeckers	307
Fig. 9.8	In the cream-coloured woodpecker, only the male bears the red beard spot	307
Fig. 9.9	The masked crimson tanager has predominantly red plumage	308
Fig. 9.10	In the pied tamarins, the alpha female mates—most unusually—with several males	309
Fig. 9.11	The large eyes of the Spix's night monkey reveal that this species is nocturnal	311
Fig. 9.12	The Brazilian porcupine is an arboreal inhabitant of the <i>terra firme</i> . The brush-tailed rat (smaller picture) is not related to the rats, but belongs to the New world porcupines, which form a separate family	312
Fig. 9.13	The golden-rumped aguti is the only animal in the rainforest capable of opening the Brazil nut	313
Fig. 9.14	The female tarantula has pink foot ends. Newly hatched tarantulas still carry a supply of yolk with them	315
Fig. 9.15	Leafcutter ants cut off pieces of leaves with which they feed the fungi grown in their burrow. This is a roundabout way for them to obtain a diet rich in protein and vitamins. Clearly visible in the upper picture are the entrances to the underground burrows of the leafcutter ants	318
Fig. 9.16	Hollow <i>internodes</i> of cecropial shoots are colonized by Azteca ants	320
Fig. 9.17	The 3 cm bullet ants are very defensible with their poisonous spines	321
Fig. 9.18	Termite nests are always built in the treetops in the rainforest. The termite queen has a shapeless swollen abdomen due to the huge number of constantly newly formed eggs	322
Fig. 9.19	The wingspan of the male giant rhinoceros beetle is 30 cm	324
Fig. 9.20	The praying mantis is a lurking hunter that waits until a suitable victim passes by	325

# List of Tables

Table 2.1	The Amazon in figures	7
Table 2.2	Geological, chemical and landscape comparison of the three Amazonian water body types	25
Table 3.1	Names of Amazon fish in Brazilian	37
Table 5.1	Development of species diversity on an alluvial area of 5000 m <sup>2</sup> (0.5 ha) in Amazonia (tree species with a diameter of at least 10 cm at breast height)	78
Table 5.2	The components of the Brazil nut—The Brazil nut with “anti-aging” properties	98
Table 8.1	Vertebrate species recorded in the Amazon lowlands and Germany	196



# 1

## Introduction: What Travellers Can Really See on Their Trip to the Rainforest

The largest river on earth has fascinated me since my student days. Since the late 1970s, I have visited the region between *Nauta* in Peru and the estuary west of *Belem* in Brazil again and again—at least 30 times to date. Initially, I also travelled to the Amazon region for scientific purposes. The main focus of my research interest at that time was the *cichlids*, a tropical fish family from the group of perch relatives (*Percomorhaceae*), which is characterized by an extremely interesting and highly developed courtship and breeding behavior. Of the approximately 3000 *cichlid species* worldwide to date, more than 500 have now been recorded in the Amazon and Rio Negro. Since the end of the 1980s, I have been leading natural history excursions into the vast landscape of the two river systems and have also been a tour guide and lecturer between *Nauta* and *Iquitos* (on the western edge of this river system) and *Belem* (about 120 river kilometres from the mouth of the Amazon into the Atlantic) on several occasions (see also the map on page II of this book).

On these journeys I had numerous encounters with a wide variety of animals as well as with members of different ethnic groups living on the banks of the rivers. My travels were also accompanied by a number of health risky adventures: there were bites from piranhas, caimans and tree snakes, attacks from swarms of bees and ant colonies, and a boat capsized on the Amazon. Fortunately, I survived all of it ultimately unscathed. To report about all this in detail would go beyond the scope of this work.

The motivation to write a summary volume on the fauna and flora of Amazonia, including the people living in this region, instead came from outside. Travel participants and listeners of my lectures have repeatedly asked for a summarizing presentation

- the fascinating flora and fauna,
- of the native peoples and the *Caboclos*, the descendants of original immigrants from Europe and these same native peoples, as well as ultimately,
- of the mighty Amazon River itself, with its enormous seasonal variations, its extremely diverse tributaries, and its general imposing characteristics

in demand. Unfortunately, this entire region is extremely endangered by constant slash and burn of the forests, by overexploitation, by illegal massive exploitation of mineral resources (e.g. gold), by the construction of huge dams, but also by the exponential growth of the human population in these sensitive ecosystems—for example, the population of Brazil has quadrupled from 1951 to 2018, from 51 million to now over 210 million. No one knows how much longer it will take until the river and its adjacent rainforests collapse. But one thing is certain: if the listed facts for the forest destruction and the river pollution and above all the exorbitant population increase do not stop by the time the next generation grows up, then in the next century there will be no more primary forests on the Amazon, and with them almost all wild animals will then also be extinct.

Perhaps the present volume will help to avert the great catastrophe that is looming in Amazonia.

I have been traveling in the Amazon lowlands for many, many years. Every time I came back, I thought on a newly started trip that I had now seen everything and there was nothing new for me to experience. But then on each subsequent trip I was surprised anew that—in addition to meeting foreign people—I kept encountering flowering plants and animals that I had never seen with my own eyes before. That's why every trip to the Amazon is full of surprises!

From the immense wealth of plant and animal species that are still far from being comprehensively identified, this book filters out, in characteristic photographs and exciting descriptions, those that visitors can actually see on a 3-week journey through the area. Each of the plant and animal examples selected in this volume has an astonishing, often unbelievable peculiarity—either in its special morphology or in its behaviours, which make you want to read or think further about the featured plant or animal.

In my research into the classification of certain plants and animals, I have discovered that there is currently a huge correction or reclassification of the long established *nomenclature*, the internationally binding scientific list of names of living things. New methods in the study of phylogeny and from molecular biology make it possible to analyse the genetic material—the genome—of a particular living being in detail. In the process, numerous new



relationships are currently being established and rearrangements as well as new species definitions are being made, which are also taken into account here.

However, the focus of this book is, as mentioned, on those plants and animals that the visitor can actually see during a flying visit. The large mammals such as the jaguar, the puma, the ocelot or the lowland tapir are therefore left unmentioned. One would first have to penetrate deep into the jungle and also live there for a longer time in order to have the chance to meet them at all. Nevertheless, the traveller will come across their traces more often, e.g. at markets where their skins are offered or tapir meat is on the menu, for example in *Iquitos*.

Dive now with me into the fascinating world of the Amazon from *Nauta* to the mouth (please see frontispiece).




# 2

## The Amazon: The Largest River on Earth

The Amazon is a river of extremes. In this chapter, its statistical data are described in detail. The reader should keep in mind, however, that all these data are provisional or approximate, as the data situation for this huge area is still incomplete today. Furthermore, the course of the river is traced from its source via various tributaries to its mouth. In the course of the last 25 million years, the geological development of this river has undergone an incredible turnaround, which can still be seen in many places on its banks. In addition, a large number of the creatures found in the river today document this reversal from a marine bay to a freshwater river. Due to the seasonal fluctuations in the river level, the morphology of both sides of the river bank is constantly changing. Both on the so-called river cliff, the steep bank on the outer side of a river bend, and on the slip-off slope, the bank on the opposite inner side, new landscape forms and river courses have developed over the years (see also Sect. 2.4). In addition, typical habitats develop during the rainy season: the flood meadows (*Várzeas*) and the flood forests (*Igapós*). These and the dry high forest (*terra firme*) are home to specific fauna and flora. Finally, the Amazon river system includes rivers with completely different chemical compositions. Thus, there are white-water, black-water and clear-water rivers, which allow the existence of very different species of plants and animals.

The largest river on our planet, with its approximately 10,000 direct or indirect tributaries and adjacent rainforests with an original size of 7,500,000 km<sup>2</sup>, constitutes the huge Amazon ecosystem directly on the equator. Figure 2.1 shows how extensive this area is: The Amazon basin including all rainforests can easily fit into the land area of the United States (without Alaska and Hawaii) with its approximately 8,000,000 km<sup>2</sup>. There are certainly



**Fig. 2.1** Size comparison of Amazonia and the land area of the United States (without Alaska and Hawaii)—Amazon basin is marked in green colour

still many unknown animal and plant species in this vast area, which has still not been fully explored. To date, more than 5500 vertebrate species and more than 56,000 seed plant species have been described in Brazil alone (for comparison, in Germany: 530 vertebrate species and about 3500 seed plant species)—and every year numerous newly discovered animal and plant species are added. The Amazon basin is already ranked first in the world for plant, mammal, fish and insect species. Probably all the species of the Amazon lowlands will never be discovered.

Among the indigenous people, the Indians, even today every few years are discovered completely isolated groups living in the rainforest without contact with modern civilization. For example, newspapers recently reported that in the Brazilian Amazon, indigenous groups spotted from a helicopter shot at the explorers' small plane with bows and arrows. According to estimates by the Brazilian Agency for the Protection of Indigenous Peoples (FUNAI), there are over 100 isolated indigenous peoples living in the Amazon rainforest alone, whose language and way of life are completely unknown.

This is all possible despite our technologically advanced and connected world, because the Amazon rainforest is so incredibly vast and yet so difficult to access. In addition, the tropical climate is extremely hostile to humans.

## 2.1 The River Between Its Source and Mouth

Since GPS calculations by Brazilian scientists in 2012, the Amazon River, which flows continuously through the South American continent from west to east at the height of the equator, is considered the longest river in the world at 6868 km, thus displacing the Nile with its 6695 km in second place (see also Table 2.1).

The source was located in 1996 in the *McIntyre Lagoon*, 200 km west of Lake Titicaca high up at 5547 m on Mount *Nevado Mismi* in the Peruvian Andes, less than 200 km from the Pacific Ocean between the cities of *Cuzco* and *Arequipa*. From there, more and more streams and rivers join together to form a common drain that flows down from this massive mountain range. This watercourse at first, bears many different names, until at last the name Amazon is found on the map: *Lloquera* at first, then *Callamayo*, *Hornilles*, *Rio Ene*, *Apurímac*, *Rio Tambo*, *Ucayali* and *Marañon* near *Nauta* in Peru, where the *Ucayali* and the *Marañon* join at the small village of *Puerto Grau*, it is called Amazonas for the first time by the Spanish-speaking people (please see page II).

There are a number of interpretations of the name “Amazonas”. It is possible, for example, that it is derived from the *Tupi-Guarini* language term “*Amaçu*”, which means “sound of water” and refers to the tidal wave of the sea, which moves upstream from the Atlantic every year in a superposition of new moon and high tide effects from two tributaries of the Amazon. Even at high tide, the water depth at *Nauta* is often only about 6 m, so even smaller

**Table 2.1** The Amazon in figures

Length <sup>a</sup>	6868 km (river kilometres)
Navigable	5215 km
Tributaries <sup>b</sup>	Approx. 10,000
Sediment transport	900 million T/year into the Atlantic Ocean
Current	0.75–5 m/s
Water volume <sup>c</sup>	
At high watermark	More than 300,000 m <sup>3</sup> /s
At low tide	About 100,000 m <sup>3</sup> /s
Gradient from the east end to the mouth	Total 180 m = 3 cm/km
Narrowest point (near Obidos)	1.5 km
Water depth	10–30 m, at Obidos deep holes of up to 120 m

<sup>a</sup>Nile: 6695 km, Rhine: 1320 km

<sup>b</sup>Of which 15 are larger than the Rhine

<sup>c</sup>In the Rhine delta 2330 km/s (max. ever measured 12,000 km/s in 1926)

cruise ships rarely venture from *Iquitos* to *Nauta*. Yet this region is highly interesting, especially from a faunal point of view. For example, within the entire Amazon lowlands, some species of the spectacularly coloured poison dart frogs (tree climbing frogs) are only found in this region, from the upper reaches to the slopes of the eastern Andes (Sect. 9.1). The rare *Gray's bald-faced saki monkey* (*Pithecia irrorata*) is also common here.

But the Amazon did not always drain in a west-east direction to the Atlantic. Before the parts of the former primeval continent *Gondwana* drifted apart, the primeval Amazon originally flowed in exactly the opposite direction from east to west and flowed into the Pacific, because about 130 million years ago there were neither the Andes nor the Atlantic. At that time, the source of the proto-Amazon was in present-day Chad (Africa) in the *Ounianga Kébir* lakes of the *Ennedi Mountains* in the middle of the Sahara (Fig. 2.2). From its source to its mouth, this river was then the longest and most powerful river on Earth at 14,000 km. With the break-up and drifting apart of the parts of the primeval continent, the Atlantic Ocean was formed and in the west of the new South American continent—beginning about 25 million years ago—the Andes gradually folded up, cutting off the estuary of the primeval Amazon from the sea and leaving many animal species of the sea in the now dammed up huge inland lake. In the course of the following many millions of years the water of this inland lake sweetened more and more, numerous former animal



Fig. 2.2 Course of the primeval Amazon about 130 million years ago

species of the sea adapted to these changed living conditions and live in the Amazon until today, for example two dolphin species, sardines, herrings, needlefish (*Belonidae*), boxfishes, rays, shrimps and crabs in more than 30 different species, sponges and many other former inhabitants of the seas. The pressure of the water masses collecting in the inland lake finally became so strong that the water now drained eastward in the same glacial valley of the primeval Amazon and lastly emptied into the Atlantic Ocean. During the two ice ages in the course of the last 100,000 years, the level of the seas sank so much that the “new” Amazon plunged over gigantic waterfalls into the Atlantic at its mouth.

There is a great deal of geological and biological evidence for the existence of the primeval Amazon and thus for the river reversal, for example:

- Rock deposits from Africa in what is now the Amazon,
- The existing anomaly that the Amazon riverbed becomes narrower towards its mouth at *Belém* (and not wider, as is usually the case with large rivers),
- Finds of various fossils, for example identical species of diatoms and *mesosaurus* in the Amazon and in the desert of Chad,
- The only desert crocodiles still living today in the lakes of the *Ennedi Mountains* (Chad), genetically related to the Amazon caimans.

By the way: even today, Sahara dust fertilizes the rainforests of the Amazon basin, as recent meteorological studies have shown. With the constant north-east trade wind, the incredible mass of 400–700 tons of it reaches every year the Amazon at an altitude of 5 km. Without this gigantic fertilizer load, the vegetation on the extremely nutrient-poor soils would never grow as lushly as we find it today.

Today, the Amazon lies in a vast lowland plain covered with sediments and sedimentary rocks, bordering the Andes to the west, the ancient low mountain ranges of the *Guyana* lands to the north, and the Brazilian Shield to the south, an ancient, eroded mountainous land. In some sections of today's Amazon, one can still see the former steep banks of the primeval Amazon: huge, million-year-old laterite cliffs rising up to 15 m, in which Amazon *Ringed kingfishers* today dig their breeding burrows (Fig. 2.3).

The slope of the river from the foot of the East Andes to its mouth is only an incredible 180 m, which is only 3 cm/km of river! Despite this extremely low gradient, the river flows at 0.75 m/s even in the dry season, which is 45 m/min or 2.7 km/h. In the rainy season, when the water masses from the many tributaries push into the main stream, the flow velocity increases up to 18 km/h. Due to the ongoing deforestation of the rainforest, the forest loses